

Finally, an internal pressure exists within a blister which varies from blister to blister. This pressure works in the opposite direction to the fluid migration into the blister space. This pressure is dependent on numerous factors, e.g., on the possibility of expansion of the blister fluid, in the slightly or severely damaged surrounding tissue, further, on the physical properties of the blister roof, i.e., thickness, degree of keratinization, elasticity etc., and thus on the location in the different body regions and the localization of the blister in the different levels of the epidermis or beneath the epidermis.

In principle it may be supposed that 2 different forces cause the fluid accumulation in blisters. On the one hand an increased filtration pressure (e.g., in urticaria bullosa or in marked inflammatory changes in the upper corium); on the other hand an increased colloid osmotic pressure in an epidermal or subepidermal cleft resulting from the skin disease initially without particularly noticeable alterations in the corium. It seems probable that both forces act in conjunction in the majority of cases of blister formation.

I am grateful to R. Brandmüller for her competent technical assistance.

#### REFERENCES

1. Braun-Falco O: The pathology of blister formation, *The Year Book of Dermatology*. Edited by AW Kopf, R Andrade, 1969, S. 6-42
2. Stoughton RB: Mechanisms of blister formation. *Arch Dermatol* 76:584-598, 1957
3. Stoughton RB: Mechanisms of vesicle formation and classification, *Dermatology in General Medicine*. Edited by RB Fitzpatrick, AA Kenneth, WH Clark, AZ Eisen, EJ Van Scott, JH Vaughan. 1971, S. 589-598
4. Stüttgen G: Zum Mechanismus der Blasenbildung an der menschlichen Haut. *Arch Dermatol Syph* 200:475-479, 1955
5. Stüttgen G: Die normale und pathologische Physiologie der Haut. Fischer, Stuttgart, 1965, S. 259
6. Kiistala U: Dermal-epidermal separation. I. The influence of age, sex and body region on suction blister formation in human skin. *Ann Clin Res* 4:10-22, 1972
7. Kiistala U, Mustakallio KK: In vivo separation of epidermis by production of suction blisters. *Lancet* I: 1444, 1964
8. Kiistala U, Mustakallio KK: Dermo-epidermal separation with suction. Electron microscopic and histo-chemical study of initial events of blistering of human skin. *J Invest Dermatol* 48:466-477, 1967
9. Granger HJ, Taylor AE, Guyton AC: Quantitative analysis of the permeability characteristics of membranes isolated from chronically implanted subcutaneous capsules. *Microvasc Res* 2:240-249, 1970
10. Guyton AC: Concept of negative interstitial pressure based on pressures in implanted perforated capsules. *Circ Res* 12:399-403, 1963
11. Guyton AC, Coleman TG: Regulation of interstitial fluid volume and pressure. *Ann NY Sci* 150:537-545, 1968
12. Guyton AC, Taylor AE, Granger HJ, Coleman TG: Interstitial fluid pressure. *Physiol Rev* 51:527-541, 1971
13. Documenta Geigy: Wissenschaftliche Tabellen. 7. Aufl. J.R. Geigy, Pharma Basel, 1969, S. 522
14. Herms W: Dehydrations- und Hyperhydrationszustände, *Pathologische Physiologie*. F Grosse-Brockhoff, Springer, Berlin-Heidelberg-New York, 1969, S. 450-456
15. Ladegaard-Pedersen HJ: Measurement of the colloid osmotic pressure in patients. *Scand J Clin Lab Invest* 20:79-87, 1967
16. Ott H: Die Errechnung des kolloidosmotischen Serumdruckes aus dem Eiweiß-Spektrum und das mittlere Molekulargewicht der Serumweißfraktionen. *Klin Wschr* 34:1079-1083, 1956
17. Guyton AC: *Textbook of Medical Physiology*. W.B. Saunders Company, Philadelphia 1976, S. 394
18. Herms W: Stoffaustausch zwischen Kapillare und Gewebe, *Pathologische Physiologie*. F Grosse-Brockhoff, Springer, Berlin-Heidelberg-New York, 1969, S. 732-736
19. Marks J, Shuster S: Disorders of capillary permeability. *Br J Dermatol* 88: 619-621, 1973

## Occupational Dermatology Symposium

An Occupational Dermatology Symposium will be held March 26-28, 1979, at the University of California Hospital, San Francisco, California. This course will succinctly summarize principles and practices of occupational dermatology, stressing the latest relevant advances. The faculty of more than 50 are international in scope. Presentations are aimed at physicians, nurses, and allied health scientists involved in the prevention and treatment of occupational skin disease. Workshops and field trips are on an optional basis. The course is jointly sponsored by the I.C.D.R.A., N.A.C.D.A., N.I.O.S.H., and the University of California. For information, write or call, Extended Programs in Medical Education, University of California Hospital, San Francisco, California, 94143, U.S.A. Telephone (415) 666-4251.